



ELSEVIER

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Linear Algebra and its Applications

journal homepage: www.elsevier.com/locate/laaOriented unicyclic graphs with the first $\left\lfloor \frac{n-9}{2} \right\rfloor$ largest skew energies[☆]Jianming Zhu^{*}*Department of Mathematics, Tongji University, Shanghai 200092, China**Business Information Management School, Shanghai Institute of Foreign Trade, Shanghai 201620, China*

ARTICLE INFO

Article history:

Received 29 February 2012

Accepted 14 June 2012

Available online 21 July 2012

Submitted by R. Brualdi

AMS classification:

05C50

15A30

Keywords:

Oriented unicyclic graphs

Skew energy

Characteristic polynomial

ABSTRACT

Let G^σ be an oriented graph obtained by assigning an orientation σ to the edge set of a simple undirected graph G such that G^σ becomes a directed graph. Let $S(G^\sigma)$ be the skew adjacency matrix of G^σ . The skew energy of G^σ is defined as the sum of the absolute values of all eigenvalues of $S(G^\sigma)$. In this paper, we provide a new method to compare the skew energies of two oriented graphs whose skew characteristic polynomials satisfy a given recurrence relation and determine the oriented unicyclic graphs of order n with the first $\left\lfloor \frac{n-9}{2} \right\rfloor$ largest skew energies for $n \geq 31$.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Let G be a simple undirected graph. The energy of G is defined as the sum of the absolute values of the eigenvalues of its adjacency matrix (see [1–3]). In theoretical chemistry, the energy of a given molecular graph is related to the total π -electron energy of the molecule represented by that graph. Consequently the graph energy has some specific chemistry interests and has been extensively studied [1–3, 12–27].

One of the graph classes that has been quite thoroughly studied is the class of all unicyclic graphs [12–24], i.e., connected graphs with one unique cycle. A number of results concerning the extremal

[☆] Supported by the National Natural Science Foundation of China No. 71173145 and Shanghai Project 085.

^{*} Address: Department of Mathematics, Tongji University, Shanghai 200092, China.

E-mail address: zhujianming2005@163.com

energies of various families of unicyclic graphs have been obtained. Hou [20] characterized the unique unicyclic graph with the minimal energy. In [16], Huo et al. determined the unique unicyclic graph with the maximal energy. Recently, Andriantiana and Wagner [15] further determined the unique unicyclic graphs with the second maximal energy.

Let G^σ be an oriented graph obtained by assigning an orientation σ to the edge set of a simple undirected graph G such that G^σ becomes a directed graph. The skew adjacency matrix $S(G^\sigma) = (s_{ij})$ of G^σ is a real skew symmetric matrix, where $s_{ij} = 1$ and $s_{ji} = -1$ if ij is an arc of G^σ with tail i and head j , otherwise $s_{ij} = s_{ji} = 0$. The skew spectrum $Sp(G^\sigma)$ of G^σ is defined as the spectrum of $S(G^\sigma)$. Note that $Sp(G^\sigma)$ consists of only purely imaginary eigenvalues because $S(G^\sigma)$ is real skew symmetric. Shader and So [7] studied the skew spectrum of oriented graphs and obtained some results.

There are various generalizations of graph energy, such as Laplacian energy, incidence energy and energy of a polynomial [4,5]. Recently, the skew energy of an oriented graph G^σ , denoted by $E_s(G^\sigma)$, is defined as the sum of the absolute values of all eigenvalues of $S(G^\sigma)$ (see [6]), that is,

$$E_s(G^\sigma) = \sum_{i=1}^n |\lambda_i|,$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ be the all eigenvalues of $S(G^\sigma)$. Recently, the skew energy of oriented graphs has been studied in [8–11].

The characteristic polynomial $\det(xI - S(G^\sigma))$ of the skew adjacency matrix $S(G^\sigma)$ of an oriented graph G^σ is also called the skew characteristic polynomial of G^σ , written as $\phi(G^\sigma, x) = \sum_{i=0}^n a_i(G^\sigma)x^{n-i}$. Since $S(G^\sigma)$ is a real skew symmetric matrix, we have $a_{2i}(G^\sigma) \geq 0$ and $a_{2i+1}(G^\sigma) = 0$ for all $0 \leq i \leq \lfloor \frac{n}{2} \rfloor$ (see [6]). Then we have

$$\phi(G^\sigma, x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{2i}(G^\sigma)x^{n-2i}. \quad (1)$$

Using these coefficients of $\phi(G^\sigma, x)$, the skew energy $E_s(G^\sigma)$ of an oriented graph G^σ of order n can be expressed by the following integral formula [11]:

$$E_s(G^\sigma) = \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \left[\sum_{i=0}^{\lfloor n/2 \rfloor} a_{2i}(G^\sigma)x^{2i} \right] dx. \quad (2)$$


Note that $a_0(G^\sigma) = 1$ and $a_2(G^\sigma)$ equals to the number of the edges of G . It follows that $E_s(G^\sigma)$ is a strictly monotonically increasing function of those numbers $a_{2i}(G^\sigma)$ ($i = 1, \dots, \lfloor \frac{n}{2} \rfloor$) for any oriented graphs. This in turn provides a way of comparing the skew energies of a pair of oriented graphs. That is to say, the method of the quasi-order relation defined by Gutman and Polansky [2] on graph energy can be generalized to the skew energy of oriented graph as follows:

Definition 1.1. Let $G_1^{\sigma_1}$ and $G_2^{\sigma_2}$ be two oriented graphs of order n . If $a_{2i}(G_1^{\sigma_1}) \leq a_{2i}(G_2^{\sigma_2})$ for all i with $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$, then we write $G_1^{\sigma_1} \preceq G_2^{\sigma_2}$.

Furthermore, if $G_1^{\sigma_1} \preceq G_2^{\sigma_2}$ and there exists at least one index j such that $a_{2j}(G_1^{\sigma_1}) < a_{2j}(G_2^{\sigma_2})$, then we write that $G_1^{\sigma_1} < G_2^{\sigma_2}$. If $a_{2i}(G_1^{\sigma_1}) = a_{2i}(G_2^{\sigma_2})$ for all i , we write $G_1^{\sigma_1} \sim G_2^{\sigma_2}$. According to the integral formula (2), we have for two oriented graphs $G_1^{\sigma_1}$ and $G_2^{\sigma_2}$ of order n that

$$\begin{aligned} G_1^{\sigma_1} \preceq G_2^{\sigma_2} &\Rightarrow E_s(G_1^{\sigma_1}) \leq E_s(G_2^{\sigma_2}); \\ G_1^{\sigma_1} < G_2^{\sigma_2} &\Rightarrow E_s(G_1^{\sigma_1}) < E_s(G_2^{\sigma_2}). \end{aligned}$$

Let G^σ be an oriented graph and C be an even cycle of G . We say C is evenly oriented relative to the orientation σ if it has an even number of edges oriented in the same direction (clockwise or anti-clockwise direction). Otherwise C is said to be oddly oriented (see Fig. 1).

Fig. 1. Two orientations of C_4 .

A linear subgraph L of G is a disjoint union of some edges and some cycles in G . We call a linear subgraph L of G evenly linear if L contains no odd cycle. We denote by $\mathcal{EL}_{2i}(G)$ the set of all evenly linear subgraphs of G with $2i$ vertices. For an evenly linear subgraph $L \in \mathcal{EL}_{2i}(G)$, we denote by $p_e(L)$ the number of evenly oriented cycles in L relative to the orientation σ .

About the coefficients $a_{2i}(G^\sigma)$ of the skew characteristic polynomial of an oriented graph G^σ , Hou and Lei [10] obtained the following result.

Lemma 1.1 [10]. *Let G^σ be an oriented graph. Then*

$$a_{2i}(G^\sigma) = \sum_{L \in \mathcal{EL}_{2i}(G)} (-1)^{p_e(L)} 2^{c(L)},$$

where $p_e(L)$ is the number of evenly oriented cycles of L and $c(L)$ is the number of even cycles of L , respectively.

In this paper, we denote by $G(n, l)$ the set of all unicyclic graphs of order n with the unique cycle C_l . A k – matching is a disjoint union of k edges in G . The number of k – matching is denoted by $m(G, k)$. We agree that $m(G, 0) = 1$ and $m(G, k) = 0$ ($k < 0$).

About the coefficients $a_{2i}(G^\sigma)$ of an oriented unicyclic graph, Hou et al. [11] obtained the following lemma.

Lemma 1.2 [11]. *Let $G \in G(n, l)$ and σ be an orientation of G . Then we have:*

- (1) If l is odd, then $a_{2i}(G^\sigma) = m(G, i)$;
- (2) If l is even and C_l is oddly oriented, then $a_{2i}(G^\sigma) = m(G, i) + 2m\left(G - C_l, i - \frac{l}{2}\right)$;
- (3) If l is even and C_l is evenly oriented, then $a_{2i}(G^\sigma) = m(G, i) - 2m\left(G - C_l, i - \frac{l}{2}\right)$.


Let $G \in G(n, l)$ and Γ be the set of all orientations of G .


Suppose that l is odd. By Lemma 1.1, $\forall \sigma_1, \sigma_2 \in \Gamma$, we have $G^{\sigma_1} \sim G^{\sigma_2}$. For convenience, $\forall \sigma \in \Gamma$, we write G^* as G^σ when l is odd.

Suppose that l is even and $\Gamma_1 = \{\sigma | \sigma \in \Gamma \text{ and } C_l \text{ is oddly oriented relative to } \sigma\}$. By Lemma 1.1, $\forall \sigma_1, \sigma_2 \in \Gamma_1$, we have $G^{\sigma_1} \sim G^{\sigma_2}$. For convenience, $\forall \sigma \in \Gamma_1$, we write G^+ as G^σ .

Suppose that l is even and $\Gamma_2 = \{\sigma | \sigma \in \Gamma \text{ and } C_l \text{ is evenly oriented relative to } \sigma\}$. By Lemma 1.2, $\forall \sigma_1, \sigma_2 \in \Gamma_2$, we have $G^{\sigma_1} \sim G^{\sigma_2}$. For convenience, $\forall \sigma \in \Gamma_2$, we write G^- as G^σ .

Let S_5^4 be the graph obtained by attaching a pendent vertex to a vertex of C_4 . We denote by $U_4(a, b)$ the graph obtained by attaching two pendent paths of length a and b to the unique pendent vertex of

Fig. 2. The graph $U_4(a, b)$ and an orientation of $U_4(a, b)$.

Fig. 3. Graphs P_n^l and F_n .

S_5^4 (see Fig. 2). When the cycle C_4 of $U_4(a, b)$ is oddly oriented relative to an orientation σ , we write $U_4^+(a, b)$ as $[U_4(a, b)]^\sigma$.

Let a, b be nonnegative integers and $\mathcal{A}_n^+ = \{U_4^+(a, b) | 0 \leq a \leq b, a + b = n - 5, a \neq 1, 3, 5\}$. In [11], Hou et al. shown that the oriented unicyclic graphs of order n with the maximal skew energy are $U_4^+(0, n - 5)$. In this paper, we further show that the oriented unicyclic graphs of order $n \geq 31$ with the first $\lfloor \frac{n-9}{2} \rfloor$ largest skew energies are the oriented unicyclic graphs in \mathcal{A}_n^+ .

The rest of this paper is organized as follows: In section 2, we investigate the skew energies of the oriented unicyclic graphs of order n where l is odd. Then section 3 is devoted to study the skew energies of the oriented unicyclic graphs of order n where l is even. In section 4, we present a new method of comparing the skew energies of two oriented graphs of order n whose skew characteristic polynomials satisfy a given recurrence relation. The oriented unicyclic graphs of order $n \geq 31$ with the first $\lfloor \frac{n-9}{2} \rfloor$ largest skew energies are determined in section 5.

2. l is odd

Let $G \in G(n, l)$ and C_l be the unique cycle of G . In this section, we always assume that l is odd.

Let P_n^l be the unicyclic graph of order n obtained by attaching a pendent path of length $n - l$ ($n > l$) to a vertex of a cycle C_l . We denote by F_n the graph obtained by attaching two pendent paths of length 2 to the unique pendent vertex of the graph P_{n-4}^4 (see Fig. 3). In this section, we will show that $G^\sigma < F_n^+$ in Lemmas 2.2 and 2.3.

The following lemma is the main method used to compare the matching numbers of two graphs in this paper.

Lemma 2.1 [4]. Let G be a graph of order n and uv be an edge of G . Then

$$m(G, k) = m(G - uv, k) + m(G - u - v, k - 1) \quad \left(1 \leq k \leq \left\lfloor \frac{n}{2} \right\rfloor\right).$$

The following result can be derived from Lemma 1.2.


Lemma 2.2. Let $G \in G(n, l)$ and σ be any orientation of G . Let l be odd. Then we have:

- (1) If n is even, then $G^\sigma < C_n^+$;
- (2) If n is odd, then $G^\sigma \preceq C_n^*$.

Proof. (1) Since l is odd, by Lemma 1.2 we have $a_{2i}(G^\sigma) = m(G, i)$. We consider the following two cases.

Case 1: $2i = n$.

By Lemma 1.2 we have, $a_n(G^\sigma) = m(G, \frac{n}{2}) \leq 2$ and $a_n(C_n^+) = m(C_n, \frac{n}{2}) + 2 = 4$. Then $a_n(G^\sigma) < a_n(C_n^+)$.

Fig. 4. The graph $P_n(a, b, c)$.

Case 2: $2i < n$.

Let uv be an edge of the unique cycle C_l of G . By Lemma 2.1 we have

$$a_{2i}(G^\sigma) = m(G, i) = m(G - uv, i) + m(G - u - v, i - 1),$$

$$a_{2i}(C_n^+) = m(C_n, i) = m(P_n, i) + m(P_{n-2}, i - 1).$$

Since $G - uv$ and $G - u - v$ are acyclic, we have $m(G - uv, i) \leq m(P_n, i)$ and $m(G - u - v, i - 1) \leq m(P_{n-2}, i - 1)$. Then $a_{2i}(G^\sigma) \leq a_{2i}(C_n^+)$. This implies that $G^\sigma < C_n^+$.

(2) The proof is similar to (1). \square

Let $P_n(a, b, c)$ be a tree of order n obtained by attaching three pendent paths of length a, b and c to an isolated vertex u where $a + b + c = n - 1$ (see Fig. 4).

Lemma 2.3. Let $n \geq 10$. Then we have:

- (1) If n is even, then $C_n^+ < F_n^+$;
- (2) If n is odd, then $C_n^* < F_n^+$.

Proof. (1) By Lemma 1.2 we have $a_{2i}(F_n^+) = m(F_n, i) + 2m(P_{n-4}(2, 2, n - 9), i - 2)$. We consider the following two cases:

Case 1: $2i = n$.

Then $a_n(F_n^+) = m(F_n, \frac{n}{2}) + 2 = 4$ and $a_n(C_n^+) = m(C_n, \frac{n}{2}) + 2 = 4$.

Case 2: $2i < n$.

Let u_3u_4 be an edge of the unique cycle C_4 of F_n (see Fig. 3). By Lemma 2.1 we have

$$\begin{aligned} m(F_n, i) &= m(F_n - u_3u_4, i) + m(F_n - u_3 - u_4, i - 1) \\ &= m(P_n(2, 2, n - 5), i) + m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1). \end{aligned}$$

Furthermore, by Lemma 2.1 we have

$$\begin{aligned} m(P_n(2, 2, n - 5), i) &= m(P_2 \cup P_{n-2}, i) + m(P_2 \cup P_{n-5}, i - 1), \\ m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1) &= m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + m(P_2 \cup P_2 \cup P_{n-9}, i - 2). \end{aligned}$$

By Lemma 1.2, we have $a_{2i}(C_n^+) = m(C_n, i) = m(P_n, i) + m(P_{n-2}, i-1)$. Moreover, by Lemma 2.1 we have

$$\begin{aligned} m(P_n, i) &= m(P_2 \cup P_{n-2}, i) + m(P_{n-3}, i-1) \\ &= m(P_2 \cup P_{n-2}, i) + m(P_2 \cup P_{n-5}, i-1) + m(P_{n-6}, i-2), \\ m(P_{n-2}, i-1) &= m(P_2 \cup P_{n-4}, i-1) + m(P_{n-5}, i-2) \\ &= m(P_2 \cup P_2 \cup P_{n-6}, i-1) + m(P_2 \cup P_{n-7}, i-2) + m(P_{n-5}, i-2). \end{aligned}$$

It follows that

$$\begin{aligned} a_{2i}(F_n^+) - a_{2i}(C_n^+) &= 2m(P_{n-4}(2, 2, n-9), i-2) + m(P_2 \cup P_2 \cup P_{n-9}, i-2) \\ &\quad - m(P_{n-6}, i-2) - m(P_2 \cup P_{n-7}, i-2) - m(P_{n-5}, i-2). \end{aligned}$$

Next,

$$\begin{aligned} m(P_{n-4}(2, 2, n-9), i-2) &= m(P_2 \cup P_{n-6}, i-2) + m(P_2 \cup P_{n-9}, i-3) \\ &= m(P_{n-6}, i-2) + m(P_{n-6}, i-3) + m(P_2 \cup P_{n-9}, i-3), \\ m(P_2 \cup P_{n-7}, i-2) - m(P_2 \cup P_2 \cup P_{n-9}, i-2) &= m(P_2 \cup P_{n-10}, i-3). \end{aligned}$$

By Lemma 2.1 we have

$$\begin{aligned} a_{2i}(F_n^+) - a_{2i}(C_n^+) &\geq m(P_{n-6}, i-2) + 2m(P_{n-6}, i-3) - m(P_{n-5}, i-2) \\ &= 2m(P_{n-6}, i-3) - m(P_{n-7}, i-3) \\ &\geq m(P_{n-6}, i-3) \\ &\geq 0. \end{aligned}$$

Then $a_{2i}(F_n^+) \geq a_{2i}(C_n^+)$. Moreover, $a_6(F_n^+) - a_6(C_n^+) \geq 1 > 0$. This implies that $C_n^+ \prec F_n^+$.
(2) The proof is similar to (1). \square

3. l is even

Let $G \in G(n, l)$ and C_l be the unique cycle of G . For convenience, we always assume that l is even in this section. The main result in this section is that if $G^\sigma \neq U_4^+(a, b)$ with $a + b = n - 5$, then $G^\sigma \preceq F_n^+$.

First, when C_l is evenly oriented, we can easily obtain the following lemma.

Lemma 3.1. *Let $G \in G(n, l)$ and l be even. Then $G^- \prec F_n^+$.*

Proof. By Lemma 1.2, we have that $a_{2i}(G^-) = m(G, i) - 2m(G - C_l, i - \frac{l}{2}) \leq m(G, i)$. Using the same method as in Lemma 2.2, we can show $G^- \preceq C_n^*$ or $G^- \preceq C_n^+$. By Lemma 2.3, the result holds. \square

Next, we need to consider the case that C_l is oddly oriented. In order to prove the main result, we take two steps to consider the problem: $l = 4$ and $l \geq 6$.

We first consider the case $l = 4$. It is easy to see that each unicyclic graph can be obtained by attaching rooted trees to the vertices of a cycle C_l . Let $G \in G(n, 4)$ and $C_4 = x_1x_2x_3x_4x_1$ be the unique cycle of G . Let $T_{a+1}, T_{b+1}, T_{c+1}, T_{d+1}$ be four rooted trees of order $a+1, b+1, c+1, d+1$, respectively. We denote by $C_4(T_a, T_b, T_c, T_d)$ the graph obtained by attaching four rooted trees $T_{a+1}, T_{b+1}, T_{c+1}, T_{d+1}$




Fig. 5. Graphs $C_4(T_a, T_b, T_c, T_d)$, $C_4(2, n-6, 0, 0)$ and $C_4(2, 0, n-6, 0)$.

to the vertex x_i ($i=1,2,3,4$) respectively. In particular, when $T_{a+1}, T_{b+1}, T_{c+1}, T_{d+1}$ are four paths with their endpoints as their roots, we write $C_4(a, b, c, d)$ as $C_4(T_a, T_b, T_c, T_d)$ (see Fig. 5).

In [25], Gutman shown that the first and second maximal energy trees of order n are P_n and $P_n(2, 2, n-5)$. The following lemma is an alternative form of Proposition 9 in [25].

Lemma 3.2 [25]. *Let T be a tree of order n . If $T \neq P_n$, then $m(T, i) \leq m(P_n(2, 2, n-5), i)$.*

The following two lemmas only are used in Theorem 3.1 to prove that if $C_4(T_a, T_b, T_c, T_d) \neq U_4(s, t)$ with $a + b + c + d = n-4$ and $s + t = n-5$, then $C_4^+(T_a, T_b, T_c, T_d) \leq F_n^+$.

Lemma 3.3. *Let $n \geq 10$. Then $C_4^+(2, 0, n-6, 0) \leq C_4^+(2, n-6, 0, 0)$.*

Proof. By Lemma 1.2 we have

$$\begin{aligned} a_{2i}(C_4^+(2, n-6, 0, 0)) &= m(C_4(2, n-6, 0, 0), i) + 2m(P_2 \cup P_{n-6}, i-2), \\ a_{2i}(C_4^+(2, 0, n-6, 0)) &= m(C_4(2, 0, n-6, 0), i) + 2m(P_2 \cup P_{n-6}, i-2). \end{aligned}$$

Let v_3v_4 be an edge of C_4 of $C_4(2, n-6, 0, 0)$ and w_3w_4 be an edge of C_4 of $C_4(2, 0, n-6, 0)$ (see Fig. 5). By Lemma 2.1, we have

$$\begin{aligned} m(C_4(2, n-6, 0, 0), i) &= m(C_4(2, n-6, 0, 0) - v_3v_4, i) + m(C_4(2, n-6, 0, 0) - v_3 - v_4, i-1) \\ &= m(P_n(2, 2, n-5), i) + m(P_4 \cup P_{n-6}, i-1), \\ m(C_4(2, 0, n-6, 0), i) &= m(C_4(2, 0, n-6, 0) - w_3w_4, i) + m(C_4(2, 0, n-6, 0) - w_3 - w_4, i-1) \\ &= m(P_n(1, 2, n-4), i) + m(P_4 \cup P_{n-6}, i-1). \end{aligned}$$

By Lemma 3.2 we have $m(P_n(2, 2, n-5), i) \geq m(P_n(1, 2, n-4), i)$. It follows that $a_{2i}(C_4^+(2, n-6, 0, 0)) \geq a_{2i}(C_4^+(2, 0, n-6, 0))$. It implies that $C_4^+(2, 0, n-6, 0) \leq C_4^+(2, n-6, 0, 0)$. \square

Lemma 3.4. *Let $n \geq 10$. Then $C_4^+(2, n-6, 0, 0) < F_n^+$.*

Proof. By Lemma 1.2, we have

$$\begin{aligned} a_{2i}(F_n^+) &= m(F_n, i) + 2m(P_{n-4}(2, 2, n-9), i-2), \\ a_{2i}(C_4^+(2, n-6, 0, 0)) &= m(C_4(2, n-6, 0, 0), i) + 2m(P_2 \cup P_{n-6}, i-2). \end{aligned}$$

Let u_3u_4 be an edge of the unique cycle C_4 of F_n and v_3v_4 be an edge of the unique cycle C_4 of $C_4(2, n-6, 0, 0)$ (see Figs. 3 and 5). By Lemma 2.1 we have

$$\begin{aligned} m(F_n, i) &= m(F_n - u_3u_4, i) + m(F_n - u_3 - u_4, i-1) \\ &= m(P_n(2, 2, n-5), i) + m(P_2 \cup P_{n-4}(2, 2, n-9), i-1), \end{aligned}$$

$$\begin{aligned} m(C_4(2, n-6, 0, 0), i) &= m(C_4(2, n-6, 0, 0) - v_3v_4, i) \\ &\quad + m(C_4(2, n-6, 0, 0) - v_3 - v_4, i-1) \\ &= m(P_n(2, 2, n-5), i) + m(P_4 \cup P_{n-6}, i-1), \end{aligned}$$

$$\begin{aligned} m(P_2 \cup P_{n-4}(2, 2, n-9), i-1) &= m(P_2 \cup P_2 \cup P_{n-6}, i-1) + m(P_2 \cup P_2 \cup P_{n-9}, i-2), \\ m(P_4 \cup P_{n-6}, i-1) &= m(P_2 \cup P_2 \cup P_{n-6}, i-1) + m(P_{n-6}, i-2). \end{aligned}$$

Thus, $m(F_n, i) - m(C_4(2, n-6, 0, 0), i) = m(P_2 \cup P_2 \cup P_{n-9}, i-2) - m(P_{n-6}, i-2)$. By Lemma 2.1 we have

$$m(P_{n-4}(2, 2, n-9), i-2) = m(P_2 \cup P_{n-6}, i-2) + m(P_2 \cup P_{n-9}, i-3).$$

It follows that

$$a_{2i}(F_n^+) - a_{2i}(C_4^+(2, n-6, 0, 0)) = 2m(P_2 \cup P_{n-9}, i-3) + m(P_2 \cup P_2 \cup P_{n-9}, i-2) - m(P_{n-6}, i-2)$$

Next,

$$\begin{aligned} m(P_{n-6}, i-2) &= m(P_3 \cup P_{n-9}, i-2) + m(P_2 \cup P_{n-10}, i-3), \\ m(P_2 \cup P_2 \cup P_{n-9}, i-2) &\geq m(P_3 \cup P_{n-9}, i-2), \\ m(P_2 \cup P_{n-9}, i-3) &\geq m(P_2 \cup P_{n-10}, i-3). \end{aligned}$$

Then $a_{2i}(F_n^+) - a_{2i}(C_4^+(2, n-6, 0, 0)) \geq m(P_2 \cup P_{n-9}, i-3) \geq 0$. Moreover, $a_6(F_n^+) - a_6(C_4^+(2, n-6, 0, 0)) \geq 1 > 0$. It implies that $C_4^+(2, n-6, 0, 0) < F_n^+$. \square

The following lemma is an alternative form of Theorem 2.2 in [26] which will be used to compare the matching numbers of two trees in Theorem 3.1 and Lemmas 3.8, 3.9.

Lemma 3.5 [26]. *Let $a + b = c + d$ with $0 \leq a \leq b$ and $0 \leq c \leq d$. Let $a < c$. Then we have:*

- (1) *If a is even, then $m(P_a \cup P_b, i) \geq m(P_c \cup P_d, i)$. Furthermore, there exists at least one index i such that the above inequality is strict.*
- (2) *If a is odd, then $m(P_a \cup P_b, i) \leq m(P_c \cup P_d, i)$. Furthermore, there exists at least one index i such that the above inequality is strict.*

Let a, b be nonnegative integers and $\mathcal{B}_n = \{U_4(a, b) | 0 \leq a \leq b, a + b = n-5\}$. We denote by $d_G(u)$ the degree of vertex u in graph G .

Theorem 3.1. *Let $G \in \mathcal{G}(n, 4)$ and $n \geq 10$. If $G \notin \mathcal{B}_n$, then $G^+ \leq F_n^+$.*

Proof. Let $C_4 = x_1x_2x_3x_4x_1$ be the unique cycle of G . Let $N(G) = \{x_i | d_G(x_i) \geq 3, i = 1, 2, 3, 4\}$. We consider the following four cases:

Case 1: $|N(G)| = 1$.

Without loss of generality, we assume $d_G(x_1) \geq 3$. Then G must be the form of $C_4(T_{n-4}, 0, 0, 0)$. By Lemma 1.2 we have

$$a_{2i}(C_4^+(T_{n-4}, 0, 0, 0)) = m(C_4(T_{n-4}, 0, 0, 0), i) + 2m(C_4(T_{n-4}, 0, 0, 0) - C_4, i - 2),$$

$$a_{2i}(F_n^+) = m(F_n, i) + 2m(P_{n-4}(2, 2, n - 9), i - 2).$$

Since $G \notin \mathcal{B}_n$, we have $C_4(T_{n-4}, 0, 0, 0) - C_4 \neq P_{n-4}$. By Lemma 3.2 we have $m(C_4(T_{n-4}, 0, 0, 0) - C_4, i - 2) \leq m(P_{n-4}(2, 2, n - 9), i - 2)$.

Let u_3u_4 be an edge of the unique cycle C_4 of F_n (see Fig. 3). By Lemma 2.1, we have

$$\begin{aligned} m(F_n, i) &= m(F_n - u_3u_4, i) + m(F_n - u_3 - u_4, i - 1) \\ &= m(P_n(2, 2, n - 5), i) + m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1), \end{aligned}$$

$$m(C_4(T_{n-4}, 0, 0, 0), i) = m(C_4(T_{n-4}, 0, 0, 0) - x_1x_2, i) + m(C_4(T_{n-4}, 0, 0, 0) - x_1 - x_2, i - 1).$$

By Lemma 3.2, we have

$$m(C_4(T_{n-4}, 0, 0, 0) - x_1x_2, i) \leq m(P_n(2, 2, n - 5), i),$$

$$m(C_4(T_{n-4}, 0, 0, 0) - x_1 - x_2, i - 1) \leq m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1).$$

It follows that $m(C_4(T_{n-4}, 0, 0, 0), i) \leq m(F_n, i)$. Then $a_{2i}(C_4^+(T_{n-4}, 0, 0, 0)) \leq a_{2i}(F_n^+)$. It implies that $C_4^+(T_{n-4}, 0, 0, 0) \leq F_n^+$.

Case 2: $|N(G)| = 2$.

Then G must be the form of $C_4(T_a, T_b, 0, 0)$ or $C_4(T_a, 0, T_b, 0)$ with $a + b = n - 4$.

Subcase 2.1: G is the form of $C_4(T_a, T_b, 0, 0)$.

Without loss of generality, we assume $d_G(x_1) \geq 3$ and $d_G(x_2) \geq 3$. By Lemma 1.2, we have

$$\begin{aligned} a_{2i}(C_4^+(T_a, T_b, 0, 0)) &= m(C_4(T_a, T_b, 0, 0), i) + 2m(C_4(T_a, T_b, 0, 0) - C_4, i - 2) \\ &\leq m(C_4(T_a, T_b, 0, 0), i) + 2m(T_a \cup T_b, i - 2), \end{aligned}$$

$$a_{2i}(C_4^+(2, n - 6, 0, 0)) = m(C_4(2, n - 6, 0, 0), i) + 2m(P_2 \cup P_{n-6}, i - 2).$$

Let v_1v_4 be an edge of C_4 of $C_4(2, n - 6, 0, 0)$ (see Fig. 5). By Lemma 2.1, we have

$$\begin{aligned} m(C_4(2, n - 6, 0, 0), i) &= m(C_4(2, n - 6, 0, 0) - v_1v_4, i) \\ &\quad + m(C_4(2, n - 6, 0, 0) - v_1 - v_4, i - 1) \\ &= m(P_n, i) + m(P_2 \cup P_2 \cup P_{n-6}, i - 1), \end{aligned}$$

$$\begin{aligned} m(C_4(T_a, T_b, 0, 0), i) &= m(C_4(T_a, T_b, 0, 0) - x_1x_2, i) \\ &\quad + m(C_4(T_a, T_b, 0, 0) - x_1 - x_2, i - 1) \\ &\leq m(P_n, i) + m(P_2 \cup T_a \cup T_b, i - 1). \end{aligned}$$

Then $m(C_4(T_a, T_b, 0, 0) - x_1x_2, i) \leq m(P_n, i)$. Since $T_a \cup T_b$ is a unconnected forest of order $n - 4$, by Lemma 3.5 we have

$$\begin{aligned} m(T_a \cup T_b, i - 2) &\leq m(P_2 \cup P_{n-6}, i - 2), \\ m(P_2 \cup T_a \cup T_b, i - 1) &\leq m(P_2 \cup P_2 \cup P_{n-6}, i - 1). \end{aligned}$$

It follows that $a_{2i}(C_4^+(T_a, T_b, 0, 0)) \leq a_{2i}(C_4^+(2, n - 6, 0, 0))$. Then $C_4^+(T_a, T_b, 0, 0) \leq C_4^+(2, n - 6, 0, 0)$. By Lemma 3.4, we have $C_4^+(T_a, T_b, 0, 0) < F_n^+$.

Subcase 2.2: G is the form of $C_4(T_a, 0, T_b, 0)$.

Using the same method as in Subcase 2.1, we can obtain that $C_4^+(T_a, 0, T_b, 0) \leq C_4^+(2, 0, n - 6, 0)$. By Lemmas 3.3 and 3.4 we have $C_4^+(T_a, 0, T_b, 0) < F_n^+$.

Case 3: $|N(G)| = 3$.

Then G must be the form of $C_4(T_a, T_b, T_c, 0)$ with $a + b + c = n - 4$. Without loss of generality, we assume $d_G(x_1) \geq 3$, $d_G(x_2) \geq 3$ and $d_G(x_3) \geq 3$. By Lemmas 1.2, 2.1, 3.5, we have

$$\begin{aligned} a_{2i}(C_4^+(T_a, T_b, T_c, 0)) &= m(C_4(T_a, T_b, T_c, 0), i) + 2m(C_4(T_a, T_b, T_c, 0) - C_4, i - 2) \\ &\leq m(C_4(T_a, T_b, T_c, 0) - x_1x_2, i) + m(C_4(T_a, T_b, T_c, 0) - x_1 - x_2, i - 1) \\ &\quad + 2m(T_a \cup T_b \cup T_c, i - 2) \\ &\leq m(P_n, i) + m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + 2m(P_2 \cup P_{n-6}, i - 2). \end{aligned}$$

According to the computing in Subcase 2.1, we have

$$a_{2i}(C_4^+(2, n - 6, 0, 0)) = m(P_n, i) + m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + 2m(P_2 \cup P_{n-6}, i - 2).$$

Then $a_{2i}(C_4^+(T_a, T_b, T_c, 0)) \leq a_{2i}(C_4^+(2, n - 6, 0, 0))$. It implies that $C_4^+(T_a, T_b, T_c, 0) \leq C_4^+(2, n - 6, 0, 0)$. By Lemma 3.4 we have $C_4^+(T_a, T_b, T_c, 0) < F_n^+$.

Case 4: $|N(G)| = 4$.

Then G must be the form of $C_4(T_a, T_b, T_c, T_d)$ with $a + b + c + d = n - 4$. By Lemmas 1.2, 2.1, 3.5, we have

$$\begin{aligned} a_{2i}(C_4^+(T_a, T_b, T_c, T_d)) &\leq m(C_4(T_a, T_b, T_c, T_d), i) + 2m(T_a \cup T_b \cup T_c \cup T_d, i - 2) \\ &= m(C_4(T_a, T_b, T_c, T_d) - x_1x_2, i) + m(C_4(T_a, T_b, T_c, T_d) \\ &\quad - x_1 - x_2, i - 1) + 2m(T_a \cup T_b \cup T_c \cup T_d, i - 2) \\ &\leq m(P_n, i) + m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + 2m(P_2 \cup P_{n-6}, i - 2). \end{aligned}$$

Then $a_{2i}(C_4^+(T_a, T_b, T_c, T_d)) \leq a_{2i}(C_4^+(2, n - 6, 0, 0))$. It follows that $C_4^+(T_a, T_b, T_c, T_d) \leq C_4^+(2, n - 6, 0, 0)$. By Lemma 3.4 we have $C_4^+(T_a, T_b, T_c, T_d) < F_n^+$.

To conclude, the result holds. \square

Next, we are ready to consider the case that $l \geq 6$. First, we will prove that if $G \neq P_n^{n-2}$, then $G^+ \leq (P_n^6)^+$. Then we will show that $(P_n^{n-2})^+ < F_n^+$ and $(P_n^6)^+ < F_n^+$.

In [10], Hou et al. proved the following result by induction on $n - l$.

Lemma 3.6 [10]. Let $G \in G(n, l)$. If $G \neq P_n^l$, then $G^+ < (P_n^l)^+$.

The following lemma about the matching number will be used in Lemmas 3.8 and 3.9 to study the skew energies of the oriented unicyclic graphs $(P_n^l)^+$.

Lemma 3.7. *Let k, r, i be nonnegative integers. If $k \geq 2r + 1$ and $i \geq r$, then $m(P_{k-2r}, i - r) \leq m(P_k, i)$.*

Proof. By Lemma 2.1 we have

$$\begin{aligned} m(P_k, i) &= m(P_{2r} \cup P_{k-2r}, i) + m(P_{2r-1} \cup P_{k-2r-1}, i - 1) \\ &= \sum_{t=0}^i [m(P_{2r}, t) \cdot m(P_{k-2r}, i - t)] + m(P_{2r-1} \cup P_{k-2r-1}, i - 1) \\ &\geq m(P_{k-2r}, i - r) + m(P_{2r-1} \cup P_{k-2r-1}, i - 1) \\ &\geq m(P_{k-2r}, i - r). \quad \square \end{aligned}$$

When n and l ($l \geq 6$) are even, the following two lemmas show that if $P_n^l \neq P_n^{n-2}$, then $(P_n^l)^+ \leq (P_n^6)^+$.

Lemma 3.8. *Let n and l be even. Then we have:*

- (1) *If $6 \leq l \leq \frac{n}{2}$, then $(P_n^l)^+ > (P_n^{l+2})^+$.*
- (2) *If $6 \leq l < \frac{n}{2} + 1$, then $(P_n^l)^+ > (P_n^{n-l+2})^+$.*

Proof.

(1) By Lemmas 1.2 and 2.1, we have

$$\begin{aligned} a_{2i}((P_n^l)^+) &= m(P_n^l, i) + 2m\left(P_{n-l}, i - \frac{l}{2}\right) \\ &= m(P_n, i) + m(P_{l-2} \cup P_{n-l}, i - 1) + 2m\left(P_{n-l}, i - \frac{l}{2}\right), \\ a_{2i}((P_n^{l+2})^+) &= m(P_n^{l+2}, i) + 2m\left(P_{n-l-2}, i - \frac{l+2}{2}\right) \\ &= m(P_n, i) + m(P_l \cup P_{n-l-2}, i - 1) + 2m\left(P_{n-l-2}, i - \frac{l+2}{2}\right). \end{aligned}$$

By Lemma 3.7 we have $m\left(P_{n-l}, i - \frac{l}{2}\right) \geq m\left(P_{n-l-2}, i - \frac{l+2}{2}\right)$.

Since $l \leq \frac{n}{2}$, we have $l - 2 \leq n - l - 2$. Since $l - 2$ is even, by Lemma 3.5 (1) we have $m(P_{l-2} \cup P_{n-l}, i - 1) \geq m(P_l \cup P_{n-l-2}, i - 1)$. Then $a_{2i}((P_n^l)^+) \geq a_{2i}((P_n^{l+2})^+)$.

When $i = \frac{l+2}{2}$, we have

$$m\left(P_{n-l}, i - \frac{l}{2}\right) = m(P_{n-l}, 1) = n - l > m\left(P_{n-l-2}, i - \frac{l+2}{2}\right) = m(P_{n-l-2}, 0) = 1.$$

Thus $a_{l+2}((P_n^l)^+) > a_{l+2}((P_n^{l+2})^+)$. It follows that $(P_n^l)^+ > (P_n^{l+2})^+$.

(2) The proof is similar to (1). \square

Lemma 3.9. *Let n be odd and $6 \leq l \leq n - 3$. If l is even, then $(P_n^l)^+ > (P_n^{l+2})^+$.*

Proof. By Lemmas 1.2 and 2.1, we have

$$\begin{aligned} a_{2i}((P_n^l)^+) &= m(P_n^l, i) + 2m\left(P_{n-l}, i - \frac{l}{2}\right) \\ &= m(P_n, i) + m(P_{l-2} \cup P_{n-l}, i - 1) + 2m\left(P_{n-l}, i - \frac{l}{2}\right), \\ a_{2i}((P_n^{l+2})^+) &= m(P_n^{l+2}, i) + 2m\left(P_{n-l-2}, i - \frac{l+2}{2}\right) \\ &= m(P_n, i) + m(P_l \cup P_{n-l-2}, i - 1) + 2m\left(P_{n-l-2}, i - \frac{l+2}{2}\right). \end{aligned}$$

By Lemma 3.7 we have $m\left(P_{n-l}, i - \frac{l}{2}\right) \geq m\left(P_{n-l-2}, i - \frac{l+2}{2}\right)$. We consider the following two cases:

Case 1: $l - 2 \leq n - l - 2$.

Since $l - 2$ is even, by Lemma 3.5 (1), we have

$$m(P_{l-2} \cup P_{n-l}, i - 1) \geq m(P_l \cup P_{n-l-2}, i - 1).$$

Thus $a_{2i}((P_n^l)^+) \geq a_{2i}((P_n^{l+2})^+)$.

Case 2: $l - 2 > n - l - 2$.

Since $n - l - 2$ is odd, by Lemma 3.5 (2), we have

$$m(P_{l-2} \cup P_{n-l}, i - 1) \geq m(P_l \cup P_{n-l-2}, i - 1).$$

Thus $a_{2i}((P_n^l)^+) \geq a_{2i}((P_n^{l+2})^+)$.

When $i = \frac{l+2}{2}$, we have

$$m\left(P_{n-l}, i - \frac{l}{2}\right) = m(P_{n-l}, 1) = n - l > m\left(P_{n-l-2}, i - \frac{l+2}{2}\right) = m(P_{n-l-2}, 0) = 1.$$

Thus $a_{l+2}((P_n^l)^+) > a_{l+2}((P_n^{l+2})^+)$. Consequently, $(P_n^l)^+ \succ (P_n^{l+2})^+$. \square

By Lemmas 3.6, 3.8 and 3.9, we can obtain that if $G \neq P_n^{n-2}, P_n^4$, then $G^+ \leq (P_n^6)^+$. In order to prove the main result in this section, we now only need to prove that $(P_n^{n-2})^+ < F_n^+$ and $(P_n^6)^+ < F_n^+$.

Lemma 3.10. If $n \geq 10$, then $(P_n^{n-2})^+ < F_n^+$.

Proof. First we prove that $(P_n^{n-2})^+ < C_n^+$. By Lemma 1.2 we have $a_n((P_n^{n-2})^+) = a_n(C_n^+) = 4$.

When $2i < n$, by Lemmas 1.2 and 2.1 we have

$$\begin{aligned} a_{2i}((P_n^{n-2})^+) &= m(P_n^{n-2}, i) + 2m(P_2, i - \frac{n-2}{2}) \\ &= m(P_n, i) + m(P_2 \cup P_{n-4}, i - 1) + 2m(P_2, i - \frac{n-2}{2}), \end{aligned}$$

$$a_{2i}(C_n^+) = m(C_n, i) = m(P_n, i) + m(P_{n-2}, i - 1).$$

We consider the following two cases:

Case 1: $2i = n - 2$.

Then

$$\begin{aligned} a_{n-2}((P_n^{n-2})^+) &= m\left(P_n, \frac{n-2}{2}\right) + m\left(P_2 \cup P_{n-4}, \frac{n-4}{2}\right) + 2, \\ a_{n-2}(C_n^+) &= m(C_n, \frac{n-2}{2}) = m\left(P_n, \frac{n-2}{2}\right) + m\left(P_{n-2}, \frac{n-4}{2}\right). \end{aligned}$$

Thus,

$$\begin{aligned} a_{n-2}(C_n^+) - a_{n-2}((P_n^{n-2})^+) &= m\left(P_{n-2}, \frac{n-4}{2}\right) - m\left(P_2 \cup P_{n-4}, \frac{n-4}{2}\right) - 2 \\ &= m\left(P_{n-5}, \frac{n-6}{2}\right) - 2 \\ &= 1 + m\left(P_{n-7}, \frac{n-8}{2}\right) - 2 \\ &= m\left(P_{n-9}, \frac{n-10}{2}\right) \\ &> 0. \end{aligned}$$

Case 2: $2i < n - 2$.

$$\begin{aligned} a_{2i}((P_n^{n-2})^+) &= m(P_n, i) + m(P_2 \cup P_{n-4}, i - 1), \\ a_{2i}(C_n^+) &= m(C_n, i) = m(P_n, i) + m(P_{n-2}, i - 1). \end{aligned}$$

By Lemma 3.5 (1), we have $a_{2i}((P_n^{n-2})^+) \leq a_{2i}(C_n^+)$. Then $(P_n^{n-2})^+ \prec C_n^+$. By Lemma 2.3 (1), we have $(P_n^{n-2})^+ \prec F_n^+$. \square

Lemma 3.11. If $n \geq 10$, then $(P_n^6)^+ \prec F_n^+$.

Proof. By Lemma 1.2 we have

$$\begin{aligned} a_{2i}((P_n^6)^+) &= m(P_n^6, i) + 2m(P_{n-6}, i - 3), \\ a_{2i}(F_n^+) &= m(F_n, i) + 2m(P_{n-4}(2, 2, n - 9), i - 2). \end{aligned}$$

Let u_3u_4 be an edge of the unique cycle C_4 of F_n (see Fig. 3). By Lemma 2.1 we have

$$\begin{aligned} m(F_n, i) &= m(F_n - u_3u_4, i) + m(F_n - u_3 - u_4, i - 1) \\ &= m(P_n(2, 2, n - 5), i) + m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1). \\ m(P_n^6, i) &= m(P_n, i) + m(P_4 \cup P_{n-6}, i - 1). \end{aligned}$$

Furthermore, by Lemma 2.1 we have

$$\begin{aligned} m(P_n(2, 2, n - 5), i) &= m(P_2 \cup P_{n-2}, i) + m(P_2 \cup P_{n-5}, i - 1), \\ m(P_n, i) &= m(P_2 \cup P_{n-2}, i) + m(P_{n-3}, i - 1) \\ &= m(P_2 \cup P_{n-2}, i) + m(P_2 \cup P_{n-5}, i - 1) + m(P_{n-6}, i - 2), \\ m(P_2 \cup P_{n-4}(2, 2, n - 9), i - 1) &= m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + m(P_2 \cup P_2 \cup P_{n-9}, i - 2), \\ m(P_4 \cup P_{n-6}, i - 1) &= m(P_2 \cup P_2 \cup P_{n-6}, i - 1) + m(P_{n-6}, i - 2). \end{aligned}$$

Then $m(F_n, i) - m(P_n^6, i) = m(P_2 \cup P_2 \cup P_{n-9}, i - 2) - 2m(P_{n-6}, i - 2)$. By Lemma 2.1 we have

$$\begin{aligned} m(P_{n-4}(2, 2, n - 9), i - 2) &= m(P_2 \cup P_{n-6}, i - 2) + m(P_2 \cup P_{n-9}, i - 3) \\ &= m(P_{n-6}, i - 2) + m(P_{n-6}, i - 3) + m(P_2 \cup P_{n-9}, i - 3). \end{aligned}$$

Thus,

$$a_{2i}(F_n^+) - a_{2i}((P_n^6)^+) = m(P_2 \cup P_2 \cup P_{n-9}, i-2) + 2m(P_2 \cup P_{n-9}, i-3) \geq 0.$$

Moreover, $a_6(F_n^+) - a_6((P_n^6)^+) = n - 6 > 0$. Then $(P_n^6)^+ \prec F_n^+$. \square

4. A new method of comparing the skew energies of two oriented graphs whose skew characteristic polynomials satisfy a given recurrence relation

In [27], Shan et al. presented a new method to compare the energies of two k -subdivision bipartite graphs. The main ideas can also apply to compare the skew energies of two oriented graphs whose skew characteristic polynomials satisfy a given recurrence relation.

In this section, we present a new method to compare the skew energies of two oriented graphs whose skew characteristic polynomials satisfy a given recurrence relation. As the applications of the new method, we can compare the skew energies of two oriented unicyclic graphs $U_4^+(7, n-12)$ and F_n^+ which are quasi-order incomparable.

Similar to the Coulson integral formula for the graph energy [1], Adiga et al. [6] obtained the integral formula for the skew energy of an oriented graph of order n as follows:

Lemma 4.1 [6]. *Let $\phi(G^\sigma, x)$ be the skew characteristic polynomial of an oriented graph G^σ of order n . Then we have*

$$E_s(G^\sigma) = \frac{1}{\pi} \int_{-\infty}^{+\infty} \left[n + x \frac{\phi'(G^\sigma, -x)}{\phi(G^\sigma, -x)} \right] dx.$$

In the following two lemmas, we show the integral formula for the difference of the skew energies of two oriented graphs of order n .

Lemma 4.2.

$$\lim_{x \rightarrow \infty} x \log \frac{x^k + a_1 x^{k-1} + \cdots + a_k}{x^k + b_1 x^{k-1} + \cdots + b_k} = a_1 - b_1.$$

Proof. Write $t = \frac{1}{x}$.

$$\begin{aligned} & \lim_{x \rightarrow \infty} x \log \frac{x^k + a_1 x^{k-1} + \cdots + a_k}{x^k + b_1 x^{k-1} + \cdots + b_k} \\ &= \lim_{t \rightarrow 0} \frac{\log(1 + a_1 t + \cdots + a_k t^k) - \log(1 + b_1 t + \cdots + b_k t^k)}{t} = a_1 - b_1. \quad \square \end{aligned}$$

Lemma 4.3. *Let $\phi(G_1^{\sigma_1}, x)$ and $\phi(G_2^{\sigma_2}, x)$ be two skew characteristic polynomials of two oriented graphs $G_1^{\sigma_1}$ and $G_2^{\sigma_2}$ of order n . Then*

$$E_s(G_1^{\sigma_1}) - E_s(G_2^{\sigma_2}) = \frac{2}{\pi} \int_0^{+\infty} \log \frac{\phi(G_1^{\sigma_1}, x)}{\phi(G_2^{\sigma_2}, x)} dx.$$

Proof. For convenience, we write $\phi_1(x) = \phi(G_1^{\sigma_1}, x)$ and $\phi_2(x) = \phi(G_2^{\sigma_2}, x)$. By Lemmas 4.1 and Lemma 4.1, we have

$$\begin{aligned}
E_s(G_1^{\sigma_1}) - E_s(G_2^{\sigma_2}) &= \frac{1}{\pi} \int_{-\infty}^{+\infty} x \left[\frac{\phi_1'(-x)}{\phi_1(-x)} - \frac{\phi_2'(-x)}{\phi_2(-x)} \right] dx \\
&= \frac{1}{\pi} \int_{-\infty}^{+\infty} x \left[\frac{\phi_1'(-x)\phi_2(-x) - \phi_1(-x)\phi_2'(-x)}{\phi_1(-x)\phi_2(-x)} \right] dx \\
&= \frac{1}{\pi} \int_{-\infty}^{+\infty} -x \left[\log \frac{\phi_1(-x)}{\phi_2(-x)} \right]' dx \\
&= -\frac{1}{\pi} x \log \frac{\phi_1(-x)}{\phi_2(-x)} \Big|_{-\infty}^{+\infty} + \frac{1}{\pi} \int_{-\infty}^{+\infty} \log \frac{\phi_1(-x)}{\phi_2(-x)} dx \\
&= \frac{1}{\pi} \int_{-\infty}^{+\infty} \log \frac{\phi_1(-x)}{\phi_2(-x)} dx \\
&= \frac{2}{\pi} \int_0^{+\infty} \log \frac{\phi_1(x)}{\phi_2(x)} dx \\
&= \frac{2}{\pi} \int_0^{+\infty} \log \frac{\phi(G_1^{\sigma_1}, x)}{\phi(G_2^{\sigma_2}, x)} dx. \quad \square
\end{aligned}$$

In the following theorem, we present a new method to comparing the skew energies of two oriented graphs whose skew characteristic polynomials satisfy a given recurrence relation.

Theorem 4.1. Let $f_n = \phi(G_1^{\sigma_1}, x)$ and $g_n = \phi(G_2^{\sigma_2}, x)$ be two skew characteristic polynomials of two oriented graphs $G_1^{\sigma_1}$ and $G_2^{\sigma_2}$ of order n . Assume that $f_n = xf_{n-1} + f_{n-2}$ and $g_n = xg_{n-1} + g_{n-2}$. Write $d_n = \frac{f_n}{g_n}$. Let k_0 and k be integers with $1 \leq k_0 \leq k \leq \lfloor \frac{n}{2} \rfloor$. Then we have:

(1) If $d_{2k_0} < d_{2k_0-1}$, then

$$E_s(G_1^{\sigma_1}) - E_s(G_2^{\sigma_2}) = \frac{2}{\pi} \int_0^{+\infty} \log d_n dx < \frac{2}{\pi} \int_0^{+\infty} \log d_{2k-1} dx.$$

(2) If $d_{2k_0} > d_{2k_0-1}$, then

$$E_s(G_1^{\sigma_1}) - E_s(G_2^{\sigma_2}) = \frac{2}{\pi} \int_0^{+\infty} \log d_n dx < \frac{2}{\pi} \int_0^{+\infty} \log d_{2k} dx.$$

Proof. (1) For convenience, we assume that $x > 0$ in what follows.

First,

$$d_{2k+2} - d_{2k} = \frac{f_{2k+2}}{g_{2k+2}} - \frac{f_{2k}}{g_{2k}} = \frac{f_{2k+2}g_{2k} - f_{2k}g_{2k+2}}{g_{2k}g_{2k+2}}.$$

Since $f_{2k+2} = xf_{2k+1} + f_{2k}$ and $g_{2k+2} = xg_{2k+1} + g_{2k}$, we have

$$\begin{aligned}
f_{2k+2}g_{2k} - f_{2k}g_{2k+2} &= x(f_{2k+1}g_{2k} - f_{2k}g_{2k+1}) \\
&= x(f_{2k-1}g_{2k} - f_{2k}g_{2k-1}) \\
&= \dots \\
&= x(f_{2k_0+1}g_{2k_0} - f_{2k_0}g_{2k_0+1}) \\
&= -x(f_{2k_0}g_{2k_0-1} - f_{2k_0-1}g_{2k_0}).
\end{aligned}$$

Then,

$$\begin{aligned} d_{2k+2} - d_{2k} &= -x \frac{f_{2k_0} g_{2k_0-1} - f_{2k_0-1} g_{2k_0}}{g_{2k} g_{2k+2}} \\ &= -x (d_{2k_0} - d_{2k_0-1}) \frac{g_{2k_0-1} g_{2k_0}}{g_{2k} g_{2k+2}} \\ &> 0. \end{aligned}$$

Similarly, we have

$$d_{2k+1} - d_{2k-1} = x (d_{2k_0} - d_{2k_0-1}) \frac{g_{2k_0-1} g_{2k_0}}{g_{2k-1} g_{2k+1}} < 0.$$

This implies that

$$d_{2k_0} < d_{2k_0+2} < \cdots < d_n < \cdots < d_{2k_0+1} < d_{2k_0-1}.$$

By Lemma 4.3, the result holds.

(2) The proof is similar to (1). \square

Next, we use Theorem 4.1 to compare the skew energies of two oriented graphs $U_4^+(7, n-12)$ and F_n^+ . Before this, we need to obtain the recurrence relation of their skew characteristic polynomials. In [10], Hou and Lei obtained the recurrence relation of the skew characteristic polynomial $\phi(G^\sigma, x)$ of an oriented graph G^σ as follows:

Lemma 4.4 [10]. *Let $e = uv$ be an edge that is on no even cycle of G and σ be an orientation of G . Then*

$$\phi(G^\sigma, x) = \phi(G^\sigma - e, x) + \phi(G^\sigma - u - v, x).$$

Let T be a tree of order n and σ_1, σ_2 be any two orientations of T . By Lemma 1.1 we have, $a_{2i}(T^{\sigma_1}) = m(T, i) = a_{2i}(T^{\sigma_2})$. Then $T^{\sigma_1} \sim T^{\sigma_2}$. For convenience, we write T^* as T^σ where σ is any orientation of T .

Lemma 4.5. *Let $n \geq 15$. Then we have:*

- (1) $\phi(F_n^+, x) = x\phi(F_{n-1}^+, x) + \phi(F_{n-2}^+, x)$.
- (2) $\phi(U_4^+(7, n-12), x) = x\phi(U_4^+(7, n-13), x) + \phi(U_4^+(7, n-14), x)$.

Proof. (1) Let u_1v_1 be a cut edge of graph F_n (see Fig. 6). By Lemma 4.4 we have

$$\phi(F_n^+, x) = \phi(F_n^+ - u_1v_1, x) + \phi(F_n^+ - u_1 - v_1, x).$$

To simplify, we let $p_n = \phi(P_n^*(2, 2, n-5), x)$. By Lemma 4.4 we have $p_n = xp_{n-1} + p_{n-2}$.




Fig. 6. Graphs F_n and $U_4(7, n-12)$.

Furthermore,

$$\begin{aligned}\phi(F_n^+ - u_1 v_1, x) &= \phi((P_5^4)^+, x) \cdot p_{n-5}, \\ \phi(F_n^+ - u_1 - v_1, x) &= \phi(C_4^+, x) \cdot p_{n-6}.\end{aligned}$$

Thus,

$$\phi(F_n^+, x) = \phi((P_5^4)^+, x) \cdot p_{n-5} + \phi(C_4^+, x) \cdot p_{n-6}.$$

Then,

$$\begin{aligned}\phi(F_n^+, x) &= (\phi((P_5^4)^+, x), \phi(C_4^+, x)) \begin{pmatrix} p_{n-5} \\ p_{n-6} \end{pmatrix} \\ &= \phi((P_5^4)^+, x), \phi(C_4^+, x) \begin{pmatrix} p_{n-6} & p_{n-7} \\ p_{n-7} & p_{n-8} \end{pmatrix} (x, 1)^T \\ &= (\phi(F_{n-1}^+, x), \phi(F_{n-2}^+, x)) \begin{pmatrix} x \\ 1 \end{pmatrix} \\ &= x\phi(F_{n-1}^+, x) + \phi(F_{n-2}^+, x).\end{aligned}$$

(2) Let $u_2 v_2$ be a cut edge of graph $U_4(7, n - 12)$ (see Fig. 6). By Lemma 4.4 we have

$$\begin{aligned}\phi(U_4^+(7, n - 12), x) &= \phi(U_4^+(7, n - 12) - u_2 v_2, x) + \phi(U_4^+(7, n - 12) - u_2 - v_2, x) \\ &= x\phi(U_4^+(7, n - 13), x) + \phi(U_4^+(7, n - 14), x). \quad \square\end{aligned}$$

Theorem 4.2. Let $n \geq 31$. Then $E_5(F_n^+) < E_5(U_4^+(7, n - 12))$.

Proof. To simplify, we let $f_n = \phi(F_n^+, x)$ and $g_n = \phi(U_4^+(7, n - 12), x)$. By Lemma 4.5 we have $f_n = xf_{n-1} + f_{n-2}$ and $g_n = xg_{n-1} + g_{n-2}$.

By some direct calculations, we have

$$\begin{aligned}f_{15} &= 26x + 181x^3 + 420x^5 + 466x^7 + 277x^9 + 90x^{11} + 15x^{13} + x^{15}, \\ g_{15} &= 24x + 180x^3 + 426x^5 + 471x^7 + 278x^9 + 90x^{11} + 15x^{13} + x^{15}, \\ f_{16} &= 4 + 96x^2 + 407x^4 + 728x^6 + 678x^8 + 354x^{10} + 104x^{12} + 16x^{14} + x^{16}, \\ g_{16} &= 4 + 96x^2 + 413x^4 + 739x^6 + 684x^8 + 355x^{10} + 104x^{12} + 16x^{14} + x^{16}.\end{aligned}$$

Furthermore,

$$f_{16}g_{15} - g_{16}f_{15} = -x(x^2 + 1)(x^{12} + 12x^{10} + 52x^8 + 97x^6 + 73x^4 + 22x^2 + 1)(x^2 + 2)^3.$$

Let $d_n = \frac{f_n}{g_n}$. If $x > 0$, then

$$d_{16} - d_{15} = \frac{f_{16}}{g_{16}} - \frac{f_{15}}{g_{15}} = \frac{f_{16}g_{15} - g_{16}f_{15}}{g_{15}g_{16}} < 0.$$

By Theorem 4.1 we have

$$E_s(F_n^+) - E_s(U_4^+(7, n-12)) = \frac{2}{\pi} \int_0^{+\infty} \log d_n dx < \frac{2}{\pi} \int_0^{+\infty} \log d_{31} dx \doteq -9.9471 \times 10^{-5} < 0.$$

Then the result holds. \square

5. Oriented unicyclic graphs of order n with the first $\lfloor \frac{n-9}{2} \rfloor$ largest skew energies

In this section, we can use the results in section 2, 3, 4 to determine the oriented unicyclic graphs of order $n \geq 31$ with the first $\lfloor \frac{n-9}{2} \rfloor$ largest skew energies in Theorem 5.2. First, we prove the following lemma.

Lemma 5.1. *Let $a + b = c + d$ with $0 \leq a \leq b$ and $0 \leq c \leq d$. Assume that $a < c$. Then we have:*

- (1) *If a is even, then $U_4^+(a, b) \succ U_4^+(c, d)$;*
- (2) *If a is odd, then $U_4^+(a, b) \prec U_4^+(c, d)$.*

Proof. By Lemma 1.2 we have

$$\begin{aligned} a_{2i}(U_4^+(a, b)) &= m(U_4(a, b), i) + 2m(P_{n-4}, i-2), \\ a_{2i}(U_4^+(c, d)) &= m(U_4(c, d), i) + 2m(P_{n-4}, i-2). \end{aligned}$$

Let vw be an edge of C_4 of graph $U_4(a, b)$ (see Fig. 2). By Lemma 2.1 we have

$$\begin{aligned} m(U_4(a, b), i) &= m(U_4(a, b) - vw, i) + m(U_4(a, b) - v - w, i-1) \\ &= m(P_n(4, a, b), i) + m(P_2 \cup P_{n-4}, i-1). \end{aligned}$$

Similarly, we have

$$m(U_4(c, d), i) = m(P_n(4, c, d), i) + m(P_2 \cup P_{n-4}, i-1).$$

Thus, $a_{2i}(U_4^+(a, b)) - a_{2i}(U_4^+(c, d)) = m(P_n(4, a, b), i) - m(P_n(4, c, d), i)$.

Moreover, by Lemma 2.1 we have

$$\begin{aligned} m(P_n(4, a, b), i) &= m(P_3 \cup P_{n-4}, i) + m(P_2 \cup P_a \cup P_b, i-1), \\ m(P_n(4, c, d), i) &= m(P_3 \cup P_{n-4}, i) + m(P_2 \cup P_c \cup P_d, i-1). \end{aligned}$$

Then $a_{2i}(U_4^+(a, b)) - a_{2i}(U_4^+(c, d)) = m(P_2 \cup P_a \cup P_b, i-1) - m(P_2 \cup P_c \cup P_d, i-1)$. Consequently, the results hold directly from Lemma 3.5. \square

Let $B_n^+ = \{U_4^+(a, b) | 0 \leq a \leq b, a + b = n - 5\}$. From Lemma 5.1, we can easily obtain the following result.

Theorem 5.1. *Let $k = \lfloor \frac{n-5}{2} \rfloor$, $t = \lfloor \frac{k}{2} \rfloor$ and $l = \lfloor \frac{k-1}{2} \rfloor$. Then we have the following quasi-order relation in B_n^+ :*

$$\begin{aligned} U_4^+(0, n-5) &\succ U_4^+(2, n-7) \succ \cdots \succ U_4^+(2t, n-5-2t) \succ U_4^+(2l+1, n-5-2l-1) \\ &\succ \cdots \succ U_4^+(7, n-12) \succ U_4^+(5, n-10) \succ U_4^+(3, n-8) \succ U_4^+(1, n-6). \end{aligned}$$

Let $\mathcal{A}_n^+ = \mathcal{B}_n^+ \setminus \{U_4^+(5, n-10), U_4^+(3, n-8), U_4^+(1, n-6)\}$.

Theorem 5.2. Let $n \geq 31$. The oriented unicyclic graphs of order n with the first $\lfloor \frac{n-9}{2} \rfloor$ largest skew energies are the oriented unicyclic graphs in \mathcal{A}_n^+ .

Proof. Let G^σ be an oriented unicyclic graph of order n and $G^\sigma \notin \mathcal{B}_n^+$. Let C_l be the unique cycle of G . By Theorem 5.1, we only need to prove that $E_s(G^\sigma) < E_s(U_4^+(7, n-12))$. We consider the following cases:

Case 1: l is odd.

By Lemmas 2.2 and 2.3, we have $G^\sigma < F_n^+$. By Theorem 4.2 we have $E_s(G^\sigma) < E_s(U_4^+(7, n-12))$.

Case 2: l is even.

By Lemma 3.1 we have $G^- < F_n^+$. By Theorem 4.2 we have $E_s(G^-) < E_s(U_4^+(7, n-12))$.

Subcase 2.1: $l = 4$.

By Theorem 3.1 we have $G^+ \leq F_n^+$. By Theorem 4.2 we have $E_s(G^+) < E_s(U_4^+(7, n-12))$.

Subcase 2.2: $l \geq 6$.

By Lemma 3.6 we have $G^+ \leq (P_n^l)^+$. By Lemmas 3.8 and 3.9, we have if $P_n^l \neq P_n^{n-2}$, then $(P_n^l)^+ \leq (P_n^6)^+$.

By Lemmas 3.10 and 3.11, we have $G^+ < F_n^+$. By Theorem 4.2 we have $E_s(G^+) < E_s(U_4^+(7, n-12))$. To conclude, the result holds. \square

Acknowledgments

The author is grateful to Professor Jia-Yu Shao for helpful discussions and his valuable suggestions about how to study the skew energy of oriented graphs. We also sincerely thank the anonymous referee for his valuable comments which greatly improve the presentation of this paper.

References

- [1] I. Gutman, The energy of a graph, Ber. Math.-Statist. Sect. Forsch. Graz. 103 (1978) 1–22.
- [2] I. Gutman, O.E. Polansky, Mathematics Concepts in Organic Chemistry, Springer-Verlag, Berlin, 1986.
- [3] I. Gutman, The Energy of a Graph: Old and New Results, Algebraic Combinatorics and Applications, Springer-Verlag, Berlin, 2001.
- [4] J. Shao, F. Gong, I. Gutman, New approaches for the real and complex integral formulas of the energy of a polynomial, Match. Commun. Math. Comput. Chem. 66 (2011) 849–861.
- [5] M. Mateljevic, V. Bozin, I. Gutman, Energy of a polynomial and the Coulson integral formula, J. Math. Chem. 48 (2010) 1062–1068.
- [6] C. Adiga, R. Balakrishnan, W. So, The skew energy of a digraph, Linear Algebra Appl. 432 (2010) 1825–1835.
- [7] B. Shader, W. So, Skew spectra of oriented graphs, Electron. J. Combin. 16 (2009) 1–6.
- [8] G. Tian, On the skew energy of orientations of hypercubes, Linear Algebra Appl. 435 (2011) 2140–2149.
- [9] S. Gong, G. Xu, 3-regular digraphs with optimum skew energy, Linear Algebra Appl. 436 (2012) 465–471.
- [10] Y. Hou, T. Lei, Characteristic polynomial of skew-adjacency matrices of oriented graphs, Electron. J. Combin. 18 (2011) P156.
- [11] Y. Hou, X. Shen, C. Zhang, Oriented unicyclic graphs with extremal skew energy, arxiv:1108.6229v1 [math.CO] 31 Aug 2011.
- [12] Y. Hou, I. Gutman, C. Woo, Unicyclic graphs with maximal energy, Linear Algebra Appl. 356 (2002) 27–36.
- [13] E.O.D. Andriantiana, S. Wagner, Unicyclic graphs with large energy, Linear Algebra Appl. 435 (2011) 1399–1414.
- [14] B. Huo, X. L. Y. Shi, Complete solution to a problem on the maximal energy of unicyclic bipartite graphs, Linear Algebra Appl. 434 (2011) 1370–1377.
- [15] E.O.D. Andriantiana, Unicyclic bipartite graphs with maximum energy, Match. Commun. Math. Comput. Chem. 66 (3) (2011) 913–926.
- [16] B. Huo, X. L. Y. Shi, Complete solution to a conjecture on the maximal energy of unicyclic graphs, Europ. J. Combin. 32 (2011) 662–673.
- [17] W. Wang, A. Chang, D. Lu, Unicyclic graphs possessing Kekule structures with minimal energy, J. Math. Chem. 42 (2007) 311–319.
- [18] W. Wang, A. Chang, L. Zhang, D. Lu, Unicyclic Huckel molecular graphs with minimal energy, J. Math. Chem. 39 (2006) 231–239.
- [19] X. L. J.B. Zhang, B. Zhou, On unicyclic conjugated molecules with minimal energies, J. Math. Chem. 42 (2007) 729–740.
- [20] Y. Hou, Unicyclic graphs with minimal energy, J. Math. Chem. 29 (3) (2001) 163–168.
- [21] Y. Liu, Some results on energy of unicyclic graphs with n vertices, J. Math. Chem. 47 (2010) 1–10.
- [22] F. Li, B. Zhou, Minimal energy of unicyclic graphs of a given diameter, J. Math. Chem. 43 (2008) 476–484.

- [23] H. Hua, M. Wang, Unicyclic graphs with given number of pendent vertices and minimal energy, *Linear Algebra Appl.* 426 (2007) 478–489.
- [24] B. Huo, J. Li, X. Li, Note on unicyclic graphs with given number of pendent vertices and minimal energy, *Linear Algebra Appl.* 433 (2010) 1381–1387.
- [25] I. Gutman, Acyclic systems with extremal Huckel π -electron energy, *Theoretical Chemistry Accounts: Theory, Computation, and Modeling (Theoret. Chim. Acta)* 45 (2) (1977) 79–87.
- [26] H. Shan, J. Shao, Graph energy change due to edge grafting operations and its applications, *MATCH Commun. Math. Comput. Chem.* 64 (2010) 25–40.
- [27] H. Shan, J. Shao, L. Zhang, C. He, A series of trees with the first $\lfloor \frac{n-7}{2} \rfloor$ largest energies, arxiv:1111.2084v1 [math. CO] 9 Nov 2011.